

Constructions of Strict Lyapunov Functions for Discrete Time and Hybrid Time-Varying Systems*

Michael Malisoff[†] Frédéric Mazenc[‡]

February 2, 2008

Abstract

We provide explicit closed form expressions for strict Lyapunov functions for time-varying discrete time systems. Our Lyapunov functions are expressed in terms of known nonstrict Lyapunov functions for the dynamics and finite sums of persistency of excitation parameters. This provides a discrete time analog of our previous continuous time Lyapunov function constructions. We also construct explicit strict Lyapunov functions for systems satisfying nonstrict discrete time analogs of the conditions from Matrosov's Theorem. We use our methods to build strict Lyapunov functions for time-varying hybrid systems that contain mixtures of continuous and discrete time evolutions.

Key Words: Strict Lyapunov functions, discrete and hybrid time-varying systems.

1 Introduction

The theory of Lyapunov functions plays a fundamental role in modern nonlinear robustness analysis and controller design [1, 3, 4, 11, 14, 15, 13, 16, 18]. In many applications, it is essential to have explicit closed form expressions for a strict Lyapunov function. This is especially the case when one wishes to design stabilizing feedbacks, which are often expressed in terms of the Lie derivatives of Lyapunov functions in the directions of the vector fields that define the system evolution. The classical converse Lyapunov function theorem asserts that systems that are stable in an appropriate sense also admit strict Lyapunov functions [5]. However, the Lyapunov functions provided by the theory are not closed form explicit expressions since they involve infinite sums or improper integrals or optimal control value functions and so do not lend themselves to applications. Moreover, whereas most of the known explicit Lyapunov function constructions are for time-invariant systems, it is well appreciated that time-invariant systems are often inadequate for engineering practice. For example, there are many applications where the dynamics cannot be stabilized by time-invariant feedback but can be stabilized using time-varying controllers [8, 19, 22, 24]. Time-varying systems are also ubiquitous in tracking. While some methods for building Lyapunov functions for time-varying systems are known, general methods for constructing *explicit closed form* Lyapunov functions for time-varying discrete and hybrid systems are not available. Hybrid systems are ubiquitous in science and engineering [27]. Hence, the construction of explicit Lyapunov functions for time-varying systems presents significant challenges that are of considerable ongoing research interest.

One recently developed and powerful approach to this problem involves constructing strict Lyapunov functions in terms of given *nonstrict* Lyapunov functions for the system; see for instance [2, 9, 16, 18]. By a nonstrict Lyapunov function, we roughly mean a function that is positive definite and radially unbounded and that has a negative semi-definite derivative along all solutions; see Section 2 for precise definitions. The advantage of this strictification approach is that in many applications, a nonstrict Lyapunov function

*Corresponding Author: Frédéric Mazenc.

[†]Department of Mathematics; Louisiana State University; Baton Rouge, LA 70803-4918; malisoff@lsu.edu. Supported by NSF Grant 0424011. The author thanks Chaohong Cai, Rafal Goebel, Ricardo Sanfelice, and Andrew Teel for illuminating discussions at the 2005 American Control Conference in Portland, Oregon.

[‡]Projet MERE INRIA-INRA; UMR Analyse des Systèmes et Biométrie INRA; 2 pl. Viala; 34060 Montpellier, France; Frederic.Mazenc@ensam.inra.fr.

is readily available through backstepping or physical considerations [13]. For continuous time systems, the strictification approach has been applied to rotating rigid bodies, robot manipulators, and other important engineering applications. This suggests the possibility of extending the strictification approach by constructing explicit closed form Lyapunov functions for discrete time time-varying systems or, more generally, hybrid time-varying systems containing both continuous and discrete time evolutions. The purpose of this work is to show that both of the extensions are indeed possible.

In Section 2, we provide the relevant definitions of strict Lyapunov functions and the necessary formalism of hybrid systems, hybrid time domains, and hybrid trajectories. In Section 3, we show how to construct explicit closed form strict Lyapunov functions for time-varying discrete time nonlinear systems in terms of given nonstrict Lyapunov functions. This provides a discrete time analog of [16] as well as a more explicit Lyapunov function construction than the known discrete time constructions that involve infinite sums of persistency of excitation (PE) parameters [20]. We also build Lyapunov functions for time-varying systems under appropriate versions of the assumptions from Matrosov's Theorem thus providing a discrete time analog of the results [18] on continuous time systems satisfying the Matrosov conditions. In Section 4, we merge our results with the known continuous time analogs to construct explicit closed form Lyapunov functions for time-varying hybrid systems, under appropriate hybrid analogs of the PE or Matrosov conditions. To our knowledge, this provides the first general method for explicitly constructing Lyapunov functions for general time-varying nonlinear hybrid systems. In Section 5, we prove our theorems. We provide some examples covered by our results in Section 6, and we close in Section 7 with some remarks about possible extensions.

2 Definitions, Assumptions, and Lemmas

We let \mathcal{K}_∞ denote the set of all continuous functions $\rho : [0, \infty) \rightarrow [0, \infty)$ for which (i) $\rho(0) = 0$ and (ii) ρ is strictly increasing and unbounded. Note that \mathcal{K}_∞ is closed under inverse and composition; i.e., if $\rho_1, \rho_2 \in \mathcal{K}_\infty$, then $\rho_1^{-1}, \rho_1 \circ \rho_2 \in \mathcal{K}_\infty$. We let \mathcal{KL} denote the class of all continuous functions $\beta : [0, \infty) \times [0, \infty) \rightarrow [0, \infty)$ for which (I) $\beta(\cdot, t) \in \mathcal{K}_\infty$ for each $t \geq 0$, (II) $\beta(s, \cdot)$ is non-increasing for each $s \geq 0$, and (III) $\beta(s, t) \rightarrow 0$ as $t \rightarrow +\infty$ for each $s \geq 0$. We let \mathcal{KLL} denote the set of all functions $\beta : [0, \infty) \times [0, \infty) \times [0, \infty) \rightarrow [0, \infty)$ such that for each $\bar{t} \geq 0$, the functions $(s, t) \mapsto \beta(s, t, \bar{t})$ and $(s, t) \mapsto \beta(s, \bar{t}, t)$ are of class \mathcal{KL} . When we say that a function ρ is *smooth* (a.k.a. C^1), we mean it is continuously differentiable, written $\rho \in C^1$. (For functions ρ defined on $[0, \infty)$, we interpret $\rho'(0)$ as a one-sided derivative, and continuity of ρ' at 0 as one-sided continuity.)

We set $\mathbb{Z}_{\geq 0} = \{0, 1, 2, \dots\}$, we let \mathbb{R}^n denote the set of all real n -tuples, and we use $|\cdot|$ to denote the usual Euclidean norm. We say that a function $\Theta : \mathbb{R}^n \times [0, \infty) \times \mathbb{Z}_{\geq 0} \rightarrow \mathbb{R} : (x, t, k) \mapsto \Theta(x, t, k)$ (which may be independent of t or k) is *uniformly state-bounded* and write $\Theta \in \mathcal{USB}$ provided there exists $\mu \in \mathcal{K}_\infty$ such that $|\Theta(x, t, k)| \leq \mu(|x|)$ for all $x \in \mathbb{R}^n$, $t \geq 0$, and $k \in \mathbb{Z}_{\geq 0}$. More generally, a vector valued function $H : \mathbb{R}^n \times [0, \infty) \times \mathbb{Z}_{\geq 0} \rightarrow \mathbb{R}^n : (x, t, k) \mapsto H(x, t, k)$ is of *class \mathcal{USB}* , written $H \in \mathcal{USB}$, provided $(x, t, k) \mapsto |H(x, t, k)|$ is of class \mathcal{USB} . Following [13], we also say Θ is *uniformly proper and positive definite (UPPD)* and write $\Theta \in \mathcal{UPPD}$ provided there are $\alpha_1, \alpha_2 \in \mathcal{K}_\infty$ such that $\alpha_1(|x|) \leq \Theta(x, t, k) \leq \alpha_2(|x|)$ for all $x \in \mathbb{R}^n$, $t \in [0, \infty)$, and $k \in \mathbb{Z}_{\geq 0}$. We say Θ is (ω_1, ω_2) -*periodic* provided $\omega_1 \in [0, \infty)$ and $\omega_2 \in \mathbb{Z}_{\geq 0}$ satisfy

$$\Theta(x, t - \omega_1, k - \omega_2) = \Theta(x, t, k) \quad \forall (x, t, k) \in \mathbb{R}^n \times [0, \infty) \times \mathbb{Z}_{\geq 0}.$$

When Θ is independent of t (resp., k), we define ω_2 -periodicity (resp., ω_1 -periodicity) analogously. A continuous function defined on a subset of Euclidean space that includes 0 and valued in $(-\infty, 0]$ is *negative semi-definite* provided it is zero at zero. A continuous function α defined on a subset of Euclidean space and valued in $[0, \infty)$ is *positive definite* provided α is zero only at zero in which case we also write $\alpha \in \mathcal{PD}$.

We study the stability properties of the discrete time fully nonlinear time-varying system

$$x_{k+1} = F(x_k, k) \tag{1}$$

where we always assume $F \in \mathcal{USB}$. We also study continuous time time-varying systems

$$\dot{x} = G(x, t) \tag{2}$$

where $G \in \mathcal{USB}$ is locally Lipschitz. We always assume (2) is *forward complete*, meaning for each $x_o \in \mathbb{R}^n$ and $t_o \geq 0$, there is a unique solution $t \mapsto \phi(t, t_o, x_o)$ for (2) defined on $[t_o, \infty)$ that satisfies $\phi(t_o, t_o, x_o) = x_o$. We

interpret the solutions of (2) in the generalized Lebesgue almost all (a.a.) sense. We also use $k \mapsto \phi(k, k_o, x_o)$ to denote the discrete time solution of (1) satisfying $\phi(k_o, k_o, x_o) = x_o$ whenever this would not lead to confusion. Given a function $V : \mathbb{R}^n \times [0, \infty) \times \mathbb{Z}_{\geq 0} \rightarrow \mathbb{R} : (x, t, k) \mapsto V(x, t, k)$, we set

$$\Delta_k V(x, t, k) := V(F(x, k), t, k + 1) - V(x, t, k), \quad \mathcal{D}V(x, t, k) := \frac{\partial V}{\partial t}(x, t, k) + \frac{\partial V}{\partial x}(x, t, k)G(x, t)$$

assuming $(x, t) \mapsto V(x, t, k)$ is also smooth for each $k \in \mathbb{Z}_{\geq 0}$ in the definition of $\mathcal{D}V$. In our analysis of (1), V will generally not depend on t but we need to allow its dependence on t in our discussion of hybrid systems.

Definition 1 (a) Let $V : \mathbb{R}^n \times \mathbb{Z}_{\geq 0} \rightarrow \mathbb{R}$ be of class $\mathcal{U}PPD$. We call V a (strict) Lyapunov function for (1) provided there exists $\alpha_3 \in \mathcal{PD}$ such that:

$$\Delta_k V(x, k) \leq -\alpha_3(|x|) \quad \forall x \in \mathbb{R}^n \ \& \ k \in \mathbb{Z}_{\geq 0}. \quad (3)$$

(b) We say that (1) is globally asymptotically stable (GAS) provided there exists $\beta \in \mathcal{KL}$ such that for all $x_o \in \mathbb{R}^n$ and $k_o \in \mathbb{Z}_{\geq 0}$, we have $|\phi(k, k_o, x_o)| \leq \beta(|x_o|, k - k_o)$ for all $k \geq k_o$.

The corresponding Lyapunov function and GAS definitions for (2) are obtained from Definition 1 by replacing k with t , $\mathbb{Z}_{\geq 0}$ with $[0, \infty)$, and $\Delta_k V$ with $\mathcal{D}V$. Notice that we do not require α_3 to be of class \mathcal{K}_∞ . For the special case where $\alpha_3 \in \mathcal{K}_\infty$, the existence of a discrete time Lyapunov function V is known to imply that (1) is GAS since then $\Delta_k V(x, k) \leq -\alpha(V(x, k))$ everywhere with $\alpha := \alpha_3 \circ \alpha_2^{-1} \in \mathcal{K}_\infty$ and α_2 as in the $\mathcal{U}PPD$ condition on V [21, Theorem 8]. Furthermore, by replacing the function $\alpha_3(|x|)$ in (3) by the smaller function $\Theta(V(x, k)) := \min\{\alpha_3(s) : \alpha_2^{-1}(V(x, k)) \leq s \leq \alpha_1^{-1}(V(x, k))\}$, we are in a situation where Lemma 15 below applies and straightforwardly implies that one can construct a Lyapunov function satisfying (3) with a new function α_3 of class \mathcal{K}_∞ . Combining this fact with the stability result from [21], we get:

Lemma 2 If (1) admits a strict Lyapunov function, then it is GAS.

We also use the following persistency of excitation (PE) notions from [13, 20]:

Definition 3 (a) We say that a bounded function $p : \mathbb{Z}_{\geq 0} \rightarrow [0, \infty)$ is of discrete PE type with parameters l and δ and write $p \in \mathcal{P}_{\text{dis}}(l, \delta)$ provided $l \in \mathbb{Z}_{\geq 0}$ and $\delta > 0$ are such that

$$\sum_{i=k-l}^k p(i) \geq \delta \quad \forall k \in \mathbb{Z}_{\geq 0}. \quad (4)$$

(b) We say that a bounded continuous function $q : [0, \infty) \rightarrow [0, \infty)$ is of continuous PE type with parameters τ and ε and write $q \in \mathcal{P}_{\text{cts}}(\tau, \varepsilon)$ provided $\tau \geq 0$ and $\varepsilon > 0$ are such that

$$\int_{t-\tau}^t q(r) dr \geq \varepsilon \quad \forall t \geq 0. \quad (5)$$

(c) We set $\mathcal{P}_{\text{dis}} = \bigcup\{\mathcal{P}_{\text{dis}}(l, \delta) : l \in \mathbb{Z}_{\geq 0}, \delta > 0\}$ and $\mathcal{P}_{\text{cts}} = \bigcup\{\mathcal{P}_{\text{cts}}(\tau, \varepsilon) : \tau \geq 0, \varepsilon > 0\}$.

Elements of \mathcal{P}_{dis} and \mathcal{P}_{cts} are called *PE parameters* and arise in a variety of contexts, e.g., $q(t) = \sin^2(t)$ as well as cases where q can be null on intervals of arbitrarily large length [13]. The following lemma follows from a simple change of variables, a Fubini Theorem argument (as was used in [13]), and the formula $1 + 2 + \dots + m = m(m + 1)/2$:

Lemma 4 Let $l \in \mathbb{Z}_{\geq 0}$, let $\tau, \varepsilon, \delta > 0$, and let $p \in \mathcal{P}_{\text{dis}}(l, \delta)$ and $q \in \mathcal{P}_{\text{cts}}(\tau, \varepsilon)$ be bounded above by \bar{p} and \bar{q} respectively. Define the functions $S : \mathbb{Z}_{\geq 0} \rightarrow [0, \infty)$ and $R : [0, \infty) \rightarrow [0, \infty)$ by

$$S(k) := \sum_{s=k-l}^k \sum_{j=s}^k p(j), \quad R(t) := \int_{t-\tau}^t \int_z^t q(\nu) d\nu dz. \quad (6)$$

Then $S(k) \leq \bar{p}(l + 1)^2$ and $R(t) \leq \tau^2 \bar{q}/2$ hold for all $k \in \mathbb{Z}_{\geq 0}$ and $t \geq 0$. If p is l -periodic, then so is S . If q is τ -periodic, then so is R .

We next recall the hybrid system tools developed in [6, 7], generalized to time-varying systems. For simplicity, we only consider singleton valued dynamics although our results carry through in the more general setting of difference and differential inclusions. Given sets $C, D \subseteq \mathbb{R}^n$ and F and G satisfying the assumptions above, the corresponding *hybrid dynamical system* is defined to be the formal object

$$\mathcal{H} := \begin{cases} \dot{x} & = G(x, t), & x \in C \\ x_{k+1} & = F(x_k, k), & x_k \in D \end{cases} . \quad (7)$$

A *compact hybrid time domain* is a subset $E \subset [0, \infty) \times \mathbb{Z}_{\geq 0}$ of the form $\cup_{k=0}^{K-1} ([t_k, t_{k+1}] \times \{k\})$ for some finite sequence $0 \leq t_0 \leq t_1 \leq \dots \leq t_K$. A *hybrid time domain* is a set $E \subset [0, \infty) \times \mathbb{Z}_{\geq 0}$ with the property that for all $(T, K) \in E$, the intersection $E \cap ([0, T] \times \{0, 1, \dots, K\})$ is a compact hybrid time domain. A *hybrid arc* is a function $x(t, k)$ defined on a hybrid time domain $\text{dom}(x)$ such that $t \mapsto x(t, k)$ is locally absolutely continuous for each k . A *hybrid trajectory* of (7) is a hybrid arc $x(t, k)$ that satisfies the following:

(S₁) For all $k \in \mathbb{Z}_{\geq 0}$ and a.a. t such that $(t, k) \in \text{dom}(x)$, we have $x(t, k) \in C$ and $\frac{\partial}{\partial t} x(t, k) = G(x(t, k), t)$.

(S₂) For all $(t, k) \in \text{dom}(x)$ such that $(t, k+1) \in \text{dom}(x)$, we have $x(t, k) \in D$ and $x(t, k+1) = F(x(t, k), k)$.

Notice that E is a hybrid time domain provided it is a finite or infinite union of sets of the form $[t_k, t_{k+1}] \times \{k\}$ with $\{t_k\}$ nondecreasing in $[0, \infty)$, with a possible additional ‘last’ set having the form $[t_k, T] \times \{k\}$ with T finite or infinite. To keep our notation simple, we use $\cup_{k \in J} ([t_k, t_{k+1}] \times \{k\})$ to denote a generic hybrid time domain with the understanding that (i) either $J = \mathbb{Z}_{\geq 0}$ or J is a finite set of the form $\{0, 1, 2, \dots, j_{\max}\}$ and (ii) $[t_k, t_{k+1}]$ may mean $[t_k, t_{k+1}]$ if J is finite and $k = j_{\max}$. Notice that continuous time solutions of (2) in C and discrete time solutions of (1) in D starting with $k = 0$ correspond to hybrid trajectories of (7) that have no switchings between the discrete and continuous evolutions.

Definition 5 (a) Let $V \in \mathcal{UPPD}$ be C^1 in x and t . We call V a (strict) Lyapunov function for \mathcal{H} provided there exists $\alpha_3 \in \mathcal{PD}$ such that the following hold for all $t \geq 0$ and $k \in \mathbb{Z}_{\geq 0}$:

$$\Delta_k V(x, t, k) \leq -\alpha_3(|x|) \quad \forall x \in D; \quad DV(x, t, k) \leq -\alpha_3(|x|) \quad \forall x \in C. \quad (8)$$

If, in addition, there is a constant $r > 0$ such that

$$V(F(x, k), t, k+1) \leq e^{-r} V(x, t, k) \quad \forall x \in D; \quad DV(x, t, k) \leq -rV(x, t, k) \quad \forall x \in C, \quad (9)$$

then we call V an exponential decay Lyapunov function for \mathcal{H} . (b) We call \mathcal{H} globally asymptotically stable (GAS) provided there exists $\beta \in \mathcal{KLL}$ such that: For each trajectory $x(t, k)$ of \mathcal{H} defined on any hybrid time domain $\cup_{k \in J} ([t_k, t_{k+1}] \times \{k\})$, we have $|x(t, k)| \leq \beta(|x(t_0, 0)|, k, t - t_k)$ for all $k \in J$ and all $t \in [t_k, t_{k+1}]$.

Lemma 6 If \mathcal{H} admits a Lyapunov function, then it is GAS.

To prove this lemma, first note that since α_3 in (8) is independent of k , standard arguments (e.g. those in [23] applied with $a(x) := \alpha_3(|x|)$) provide $\beta_1 \in \mathcal{KL}$ such that for each hybrid trajectory $x(t, j)$ defined on a hybrid time domain $\cup_{k \in J} ([t_k, t_{k+1}] \times \{k\})$ and satisfying any initial condition $x(t_0, 0) = x_0$, we have

$$|x(t, k)| \leq \beta_1(|x(t_k, k)|, t - t_k) \quad \forall k \in J, \quad t \in [t_k, t_{k+1}]. \quad (10)$$

Similarly, since α_3 in (8) is independent of t , and since we can assume as above that $\alpha_3 \in \mathcal{K}_\infty$, the argument from [21, Theorem 8] provides $\beta_2 \in \mathcal{KL}$ such that

$$|x(t_k, k)| \leq \beta_2(|x_0|, k) \quad \forall k \in J. \quad (11)$$

In fact, β_2 can be constructed using the decay conditions from (8) as follows. First note that by arguing as in the proof of Lemma 2 above and replacing V with $\kappa \circ V$ for a suitable function $\kappa \in \mathcal{K}_\infty$ in the discrete decay condition without relabeling, we can find $\gamma \in \mathcal{K}_\infty$ such that $\Delta_k V(x, t, k) \leq -\gamma(V(x, t, k))$ when $x \in D$; see Lemma 15 for the construction of κ . Since $t \mapsto V(x(t, k), t, k)$ decays on (t_k, t_{k+1}) for each k , we get $V(x(t_{k+2}, k+1), t_{k+2}, k+1) \leq V(x(t_{k+1}, k+1), t_{k+1}, k+1) = V(F(x(t_{k+1}, k), k), t_{k+1}, k+1)$ so

$$\begin{aligned} V(x(t_{k+2}, k+1), t_{k+2}, k+1) - V(x(t_{k+1}, k), t_{k+1}, k) &\leq \Delta_k V(x(t_{k+1}, k), t_{k+1}, k) \\ &\leq -\gamma(V(x(t_{k+1}, k), t_{k+1}, k)) \end{aligned}$$

everywhere. Applying [21, Theorem 8] to the function $k \mapsto V(x(t_{k+1}, k), t_{k+1}, k)$ and recalling that V is uniformly proper and positive definite gives $\tilde{\beta}_2 \in \mathcal{KL}$ (not depending on the choice of the trajectory) such that $|x(t_{k+1}, k)| \leq \tilde{\beta}_2(|x(t_1, 0)|, k)$ for all k . Choosing $\alpha_1, \alpha_2 \in \mathcal{K}_\infty$ such that $\alpha_1(|x|) \leq V(x, t, k) \leq \alpha_2(|x|)$ everywhere, the discrete time decay condition in (8) gives

$$|x(t_{k+1}, k)| \geq \alpha_2^{-1} \circ V(x(t_{k+1}, k), t_{k+1}, k) \geq \alpha_2^{-1} \circ V(x(t_{k+1}, k+1), t_{k+1}, k+1) \geq \alpha_2^{-1} \circ \alpha_1(|x(t_{k+1}, k+1)|)$$

for all $k \in J$. Similarly, the continuous time decay condition in (8) gives

$$|x(t_1, 0)| \leq \alpha_1^{-1} \circ V(x(t_1, 0), t_1, 0) \leq \alpha_1^{-1} \circ V(x(t_o, 0), t_o, 0) \leq \alpha_1^{-1} \circ \alpha_2(|x_o|).$$

We can therefore satisfy (11) by taking $\beta_2(s, k) := \alpha_1^{-1} \circ \alpha_2 \circ \tilde{\beta}_2(\alpha_1^{-1} \circ \alpha_2(s), k) + s/(k+1)$, where the additional term $s/(k+1)$ is used to account for the case $k=0$. Combining (10)-(11) shows we can satisfy the requirements of Lemma 6 using $\beta(s, t, k) = \beta_1(\beta_2(s, k), t)$.

3 Statement of Results on Discrete-Time Systems

3.1 Strictifying Persistence of Excitation (PE) Decay Estimates

We begin by constructing explicit closed form Lyapunov functions for discrete time systems in terms of nonstrict Lyapunov functions and appropriate PE parameters $p \in \mathcal{P}_{\text{dis}}$. For an alternative construction, involving infinite sums of PE parameter values, see [20]. We prove the following in Section 5:

Theorem 7 *Let $l \in \mathbb{Z}_{\geq 0}$, $\delta > 0$, $p \in \mathcal{P}_{\text{dis}}(l, \delta)$, $V \in \mathcal{UPPD}$, and $\Theta \in \mathcal{PD}$ satisfy*

$$\Delta_k V(x, k) \leq -p(k+1)\Theta(V(x, k)) \quad \forall x \in \mathbb{R}^n \ \& \ k \in \mathbb{Z}_{\geq 0}. \quad (12)$$

Then one can construct $\kappa, \gamma \in \mathcal{K}_\infty$ such that

$$U(x, k) := \kappa(V(x, k)) + \frac{\gamma(V(x, k))}{4(l+1)} \sum_{s=k-l}^k \sum_{j=s}^k p(j) \quad (13)$$

is a strict Lyapunov function for (1), so (1) is GAS. If p and V are also both l -periodic in k , then so is U .

Remark 8 *A key feature in (12) is that the PE condition on p allows $p(k+1) = 0$ for some values of k in which case we could have $\Delta_k V(x, k) = 0$. An additional novel feature of Theorem 7 is that we do not require the gain function Θ in (12) to be of class \mathcal{K}_∞ . This properness of the gain function was required in [13, 20]. Our proof of Theorem 7 will show that we can take $\kappa(s) \equiv s$ if $\Theta \in \mathcal{K}_\infty$.*

3.2 Lyapunov Function Constructions Under Matrosov Conditions

Recall the definitions of \mathcal{UPPD} and \mathcal{USB} from Section 2. We explicitly construct a Lyapunov function for discrete time systems (1) satisfying the following analog of the Matrosov Theorem conditions from [18]:

Assumption 9 *There exist $V_1 : \mathbb{R}^n \times \mathbb{Z}_{\geq 0} \rightarrow [0, \infty)$ of class \mathcal{UPPD} , $V_2 : \mathbb{R}^n \times \mathbb{Z}_{\geq 0} \rightarrow \mathbb{R}$ of class \mathcal{USB} , a function $\phi_2 \in \mathcal{K}_\infty$, nonnegative functions $N_1, N_2 \in \mathcal{USB}$, a function $\chi : \mathbb{R}^n \times [0, \infty) \times \mathbb{Z}_{\geq 0} \rightarrow \mathbb{R}$, a positive increasing function ϕ_1 , a positive definite function W , and $p \in \mathcal{P}_{\text{dis}}$ such that*

$$\Delta_k V_1(x, k) \leq -N_1(x, k),$$

$$\Delta_k V_2(x, k) \leq -N_2(x, k) + \chi(x, N_1(x, k), k),$$

$$|\chi(x, N_1(x, k), k)| \leq \phi_1(|x|)\phi_2(N_1(x, k)), \quad \text{and} \quad N_1(x, k) + N_2(x, k) \geq p(k+1)W(x)$$

hold for all $x \in \mathbb{R}^n$ and $k \in \mathbb{Z}_{\geq 0}$.

Notice that we allow V_2 to take both positive and negative values. In Section 5, we prove:

Theorem 10 *If (1) satisfies Assumption 9, then one can construct an explicit strict Lyapunov function for (1). In particular, (1) is GAS.*

4 Statement of Results on Hybrid Systems

4.1 Hybrid Persistency of Excitation Estimates

We next extend Theorem 7 to hybrid systems. To keep the exposition simple, we assume the gain functions Θ in (12) and its continuous analog are $\Theta(s) = s$, but the extension to general positive definite Θ can be done using similar arguments. We prove the following in Section 5:

Theorem 11 *Let $V \in \mathcal{U}PPD$ be C^1 in x and t . Consider the hybrid system \mathcal{H} in (7), and let $\delta, \varepsilon, \tau > 0$ and $l \in \mathbb{Z}_{\geq 0}$ be given. Assume there exist $r \in \mathcal{P}_{\text{dis}}(l, \delta)$ and $q \in \mathcal{P}_{\text{cts}}(\tau, \varepsilon)$ such that*

$$V(F(x, k), t, k+1) \leq e^{-r(k+1)}V(x, t, k) \quad \forall x \in D; \quad \mathcal{D}V(x, t, k) \leq -q(t)V(x, t, k) \quad \forall x \in C \quad (14)$$

hold for all $t \geq 0$ and $k \in \mathbb{Z}_{\geq 0}$. Then

$$V^\sharp(x, t, k) = \left[2 + \frac{1}{4(l+1)} \sum_{s=k-l}^k \sum_{j=s}^k (1 - e^{-r(j)}) + \frac{1}{\tau} \int_{t-\tau}^t \int_z^t q(\nu) d\nu dz \right] V(x, t, k) \quad (15)$$

is an exponential decay Lyapunov function for \mathcal{H} which is therefore GAS. If in addition V is (τ, l) -periodic and r and q are l -periodic and τ -periodic respectively, then V^\sharp is also (τ, l) -periodic.

Remark 12 *The preceding theorem covers continuous dynamics (by taking $D = \emptyset$ and $C = \mathbb{R}^n$ with the understanding that the term involving the double sum in V^\sharp is not present) and discrete dynamics (by taking $C = \emptyset$ and $D = \mathbb{R}^n$ in which case the term involving the double integral in V^\sharp is not present). See [6] for an alternative, nonexplicit construction of a Lyapunov function for time-invariant hybrid systems.*

4.2 Hybrid Systems Satisfying Matrosov Conditions

We next extend Theorem 10 to hybrid systems that satisfy the following analog of Assumption 9.

Assumption 13 *There exist $V_1 \in \mathcal{U}PPD$ and $V_2 \in \mathcal{U}SB$ that are C^1 in (x, t) , nonnegative $N_1, N_2 \in \mathcal{U}SB$, a function $\chi : \mathbb{R}^n \times [0, \infty)^2 \times \mathbb{Z}_{\geq 0} \rightarrow \mathbb{R}$, a positive increasing ϕ_1 , and a positive definite function W , $p \in \mathcal{P}_{\text{dis}}$, $\phi_2 \in \mathcal{K}_\infty$, and $q \in \mathcal{P}_{\text{cts}}$ such that*

1. *For all $x \in D$, we have $\Delta_k V_1(x, t, k) \leq -N_1(x, t, k)$, $\Delta_k V_2(x, t, k) \leq -N_2(x, t, k) + \chi(x, N_1(x, t, k), t, k)$, and $N_1(x, t, k) + N_2(x, t, k) \geq p(k+1)W(x)$.*
2. *For all $x \in C$, we have $\mathcal{D}V_1(x, t, k) \leq -N_1(x, t, k)$, $\mathcal{D}V_2(x, t, k) \leq -N_2(x, t, k) + \chi(x, N_1(x, t, k), t, k)$, and $N_1(x, t, k) + N_2(x, t, k) \geq q(t)W(x)$.*
3. *For all $x \in \mathbb{R}^n$, we have $|\chi(x, N_1(x, t, k), t, k)| \leq \phi_1(|x|)\phi_2(N_1(x, t, k))$.*

hold for all $t \geq 0$ and $k \in \mathbb{Z}_{\geq 0}$.

Assumption 13 simply means the discrete and continuous parts of \mathcal{H} satisfy the appropriate discrete and continuous Matrosov conditions. It reduces to Assumption 9 for discrete systems when $C = \emptyset$ and $D = \mathbb{R}^n$ in which case its condition 2. holds vacuously. Notice that we again do not require V_2 to be nonnegative. In Section 5, we prove:

Theorem 14 *If \mathcal{H} satisfies Assumption 13, then one can construct an explicit closed form strict Lyapunov function for \mathcal{H} . In particular, \mathcal{H} is GAS.*

5 Proofs of Theorems

5.1 Results on Discrete Systems

5.1.1 Proof of Theorem 7

By minorizing $\Theta \in \mathcal{P}D$ without relabeling as in [17], we assume in the sequel that $\Theta \in C^1$ is nondecreasing on $[0, 1]$ and nonincreasing on $[1, \infty)$. The next technical lemma allows us to assume that $\Theta \in \mathcal{K}_\infty$ in (12):

Lemma 15 Let $\Theta \in \mathcal{PD}$ be as above and $p \in \mathcal{P}_{\text{dis}}$. Define $\mu : [0, \infty) \rightarrow [1, \infty)$, κ , and χ by

$$\kappa(r) := 2 \int_0^r \mu(z) dz, \quad \chi(r) := \Theta(2r)\mu(r), \quad \text{and} \quad \mu(r) = \begin{cases} 1 + 4r^2, & 0 \leq r \leq 1/2 \\ \frac{4\Theta(1)r}{\Theta(2r)}, & 1/2 \leq r < \infty \end{cases}. \quad (16)$$

Let $\nu \in \mathcal{UPPD}$ satisfy $\Delta_k \nu(x, k) \leq -p(k+1)\Theta(\nu(x, k))$ for all $x \in \mathbb{R}^n$ and $k \in \mathbb{Z}_{\geq 0}$. Then $\kappa \in \mathcal{K}_{\infty} \cap C^1$, $\chi \in \mathcal{K}_{\infty}$, and $V := \kappa(\nu) \in \mathcal{UPPD}$ satisfies

$$\Delta_k V(x, k) \leq -p(k+1)\gamma(V(x, k)) \quad \forall x \in \mathbb{R}^n \ \& \ k \in \mathbb{Z}_{\geq 0}, \quad (17)$$

where $\gamma \in \mathcal{K}_{\infty}$ is defined by $\gamma(s) := \chi(\kappa^{-1}(s)/2)$.

To prove Lemma 15, fix $x \in \mathbb{R}^n$ and $k \in \mathbb{Z}_{\geq 0}$ and apply the Fundamental Theorem of Calculus to $s \mapsto \mathcal{F}(s) := \kappa(s\nu(F(x, k), k+1) + (1-s)\nu(x, k))$ to write $\Delta_k V(x, k) = \mathcal{F}(1) - \mathcal{F}(0) = \int_0^1 \mathcal{F}'(s) ds$ and so also

$$\begin{aligned} \Delta_k V(x, k) &= \left[\int_0^1 \kappa'(s\nu(F(x, k), k+1) + (1-s)\nu(x, k)) ds \right] [\nu(F(x, k), k+1) - \nu(x, k)] \\ &\leq -p(k+1) \left[\int_0^1 \kappa'(s\nu(F(x, k), k+1) + (1-s)\nu(x, k)) ds \right] \Theta(\nu(x, k)) \\ &\leq -p(k+1) \left[\int_0^1 \kappa'((1-s)\nu(x, k)) ds \right] \Theta(\nu(x, k)) \\ &\leq -p(k+1) \left[\int_0^{1/2} \kappa' \left(\frac{1}{2} \nu(x, k) \right) ds \right] \Theta(\nu(x, k)) = -p(k+1) \mu \left(\frac{1}{2} \nu(x, k) \right) \Theta(\nu(x, k)) \end{aligned}$$

where the first inequality holds because κ is nondecreasing and the other inequalities used the fact that κ' is nondecreasing. The lemma now follows from our choices of γ and χ .

We can therefore assume that V satisfies (17) with $\gamma \in \mathcal{K}_{\infty}$, possibly by replacing V with $\kappa(V)$ for $\kappa \in \mathcal{K}_{\infty}$ defined in (16). Defining $S(k)$ as in (6) and defining U by (13) with $\kappa(s) \equiv s$ therefore gives

$$\begin{aligned} \Delta_k U(x, k) &= V(F(x, k), k+1) + \frac{S(k+1)}{4(l+1)} \gamma(V(F(x, k), k+1)) - V(x, k) - \frac{S(k)}{4(l+1)} \gamma(V(x, k)) \\ &= \Delta_k V(x, k) + \frac{1}{4(l+1)} S(k+1) \Delta_k (\gamma \circ V)(x, k) + \frac{1}{4(l+1)} \gamma(V(x, k)) [S(k+1) - S(k)] \\ &\leq \Delta_k V(x, k) + \frac{1}{4(l+1)} \gamma(V(x, k)) [S(k+1) - S(k)], \end{aligned} \quad (18)$$

where the last inequality holds because γ is increasing, so $\Delta_k (\gamma \circ V)(x, k) \leq 0$. Note that

$$\begin{aligned} S(k+1) - S(k) &= \sum_{s=k+1-l}^k \sum_{j=s}^k p(j) + (l+1)p(k+1) - \sum_{s=k-l}^k \sum_{j=s}^k p(j) \\ &= \sum_{s=k-l}^k \sum_{j=s}^k p(j) - \sum_{j=k-l}^k p(j) + (l+1)p(k+1) - \sum_{s=k-l}^k \sum_{j=s}^k p(j) \\ &= - \sum_{j=k-l}^k p(j) + (l+1)p(k+1). \end{aligned} \quad (19)$$

Substituting (19) into (18) gives

$$\begin{aligned} \Delta_k U(x, k) &\leq \Delta_k V(x, k) + \frac{1}{4(l+1)} \gamma(V(x, k)) \left((l+1)p(k+1) - \sum_{j=k-l}^k p(j) \right) \\ &\leq \Delta_k V(x, k) + \frac{p(k+1)\gamma(V(x, k))}{4} - \frac{\gamma(V(x, k))}{4(l+1)} \sum_{j=k-l}^k p(j) \leq -\frac{\delta}{4(l+1)} \gamma(V(x, k)), \end{aligned} \quad (20)$$

where the last inequality follows from the PE property of p and (17). This, the fact that $V \in \mathcal{UPPD}$, and the global boundedness of $S(k)$ from Lemma 4 shows that U is a Lyapunov function for (1). Therefore, (1) is GAS, by Lemma 2. The assertion in the theorem about periodicity follows from Lemma 4 and the formula for U , so this proves our theorem.

5.1.2 Proof of Theorem 10

Let $V_3 = V_1 + V_2$, and let $\alpha_1, \alpha_2 \in \mathcal{K}_\infty$ satisfy the UPPD requirements for V_1 . In the rest of the proof, all inequalities should be interpreted as holding globally unless otherwise indicated. We also leave out the argument (x, k) of some of our \mathcal{USB} functions when this would not lead to confusion. It follows from Assumption 9 that we can determine a positive definite function λ such that

$$\Delta_k V_3(x, k) \leq -p(k+1)W(x) + \phi_1(|x|)\phi_2(N_1(x, k)) \leq -p(k+1)\lambda(V_1(x, k)) + \phi_1(|x|)\phi_2(N_1(x, k)) \quad (21)$$

e.g. $\lambda(s) = \min\{W(x) : x \in \mathbb{R}^n, \alpha_1(|x|) \leq s \leq \alpha_2(|x|)\}$ (which is positive definite because $W \in \mathcal{PD}$). By minorizing λ as necessary as in [17], we can assume it is C^1 , nondecreasing on $[0, 1/2]$ and nonincreasing on $[1/2, \infty)$. The proof of Lemma 15 above with $\Theta(r) := \lambda(r/2)$ provides an increasing continuous function $k_1 : [0, \infty) \rightarrow [1, \infty)$ such that $\Lambda_1(s) := k_1(s)\lambda(s)$ is of class \mathcal{K}_∞ . Let $V_4 = k_1(V_1)V_3$. Then (21) gives

$$\begin{aligned} \Delta_k V_4(x, k) &= [k_1(V_1(F(x, k), k+1)) - k_1(V_1(x, k))] V_3(F(x, k), k+1) \\ &\quad + k_1(V_1(x, k)) [V_3(F(x, k), k+1) - V_3(x, k)] \\ &\leq [k_1(V_1(F(x, k), k+1)) - k_1(V_1(x, k))] V_3(F(x, k), k+1) \\ &\quad - k_1(V_1(x, k))p(k+1)\lambda(V_1(x, k)) + k_1(V_1(x, k))\phi_1(|x|)\phi_2(N_1(x, k)). \end{aligned} \quad (22)$$

Since $F, V_3 \in \mathcal{USB}$ and $V_1 \in \mathcal{UPPD}$, we get continuous increasing positive functions Γ and Λ_2 such that

$$\Delta_k V_4(x, k) \leq [-\Delta_k V_1(x, k)]\Gamma(V_1(x, k)) - p(k+1)\Lambda_1(V_1(x, k)) + \Lambda_2(V_1(x, k))\phi_2(N_1(x, k)) \quad (23)$$

e.g. by first finding an increasing positive function $\tilde{\alpha}$ such that $|k'_1(r)| \leq \tilde{\alpha}(r)$. Define $k_2 \in \mathcal{K}_\infty$ by $k_2(s) = s\Gamma(s)$. Since $k_2(s)/s$ is increasing, we have $k_2(b) - k_2(a) \geq (b-a)\Gamma(b)$ when $b \geq a \geq 0$ (since $(k_2(b) - k_2(a))(b-a)^{-1} \geq k_2(b)b^{-1}$ when $b > a \geq 0$). Hence by choosing $a = V_1(F(x, k), k+1)$ and $b = V_1(x, k)$, we get $\Delta_k(k_2 \circ V_1)(x, k) \leq \Delta_k V_1(x, k)\Gamma(V_1(x, k))$ everywhere. Therefore, by adding a \mathcal{K}_∞ function to k_2 as necessary, we can assume $V_5 := V_4 + k_2(V_1) \in \mathcal{UPPD}$ and satisfies

$$\Delta_k V_5(x, k) \leq -p(k+1)\Lambda_1(V_1(x, k)) + \Lambda_2(V_1(x, k))\phi_2(N_1(x, k)). \quad (24)$$

Arguing as in the proof of Theorem 7 except with γ replaced by Λ_1 provides $V_6 \in \mathcal{UPPD}$ such that

$$\Delta_k V_6(x, k) \leq -\frac{\delta}{4(l+1)}\Lambda_1(V_1(x, k)) + \Lambda_2(V_1(x, k))\phi_2(N_1(x, k)). \quad (25)$$

By arguing as in [17, Section IV.A], we can select $k_3 \in C^1 \cap \mathcal{PD}$ so that $|k'_3(s)| \leq 1$ for all $s \geq 0$ and

$$k_3(r) \leq \phi_2^{-1} \left(\frac{\delta}{8(l+1)} \frac{\Lambda_1(r)}{1 + \Lambda_2(r)} \right) \frac{1}{1 + \Lambda_2(r)}, \text{ hence } \phi_2(k_3(V_1)\Lambda_2(V_1))\Lambda_2(V_1) \leq \frac{\delta}{8(l+1)}\Lambda_1(V_1) \quad (26)$$

everywhere. Choose $\mu_F, \alpha_6 \in \mathcal{K}_\infty$ such that $V_6(x, k) \leq \alpha_6(|x|)$ and $|F(x, k)| \leq \mu_F(|x|)$ everywhere. Arguing as above (with Γ replaced by $\alpha_6 \circ \mu_F \circ \alpha_1^{-1}$) and recalling that $\Delta_k V_1(x, k) \leq 0$ provides $k_4 \in \mathcal{K}_\infty$ such that

$$\Delta_k(k_4 \circ V_1)(x, k) \leq \alpha_6 \circ \mu_F \circ \alpha_1^{-1} \circ V_1(x, k)\Delta_k V_1(x, k) \leq \alpha_6(\mu_F(|x|))\Delta_k V_1(x, k)$$

everywhere. Since $|k'_3(s)| \leq 1$ for all $s \geq 0$, we have

$$[\Delta_k(k_3 \circ V_1)(x, k)]V_6(F(x, k), k+1) \leq [-\Delta_k V_1(x, k)]V_6(F(x, k), k+1) \leq -\alpha_6(\mu_F(|x|))\Delta_k V_1(x, k)$$

everywhere. It follows that $V_7 := k_3(V_1)V_6 + k_4(V_1) \in \mathcal{UPPD}$ satisfies

$$\begin{aligned} \Delta_k V_7(x, k) &= [\Delta_k(k_3 \circ V_1)(x, k)]V_6(F(x, k), k+1) + k_3(V_1(x, k))\Delta_k V_6(x, k) + \Delta_k(k_4 \circ V_1)(x, k) \\ &\leq -\frac{\delta}{4(l+1)}k_3(V_1)\Lambda_1(V_1) + k_3(V_1)\Lambda_2(V_1)\phi_2(N_1(x, k)). \end{aligned} \quad (27)$$

Next note that for all functions $\mu \in \mathcal{K}_\infty$,

$$\Delta_k V_7 \leq -\frac{\delta}{4(l+1)}k_3(V_1)\Lambda_1(V_1) + \mu(k_3(V_1)\Lambda_2(V_1))k_3(V_1)\Lambda_2(V_1) + \mu^{-1}(\phi_2(N_1(x, k)))\phi_2(N_1(x, k)). \quad (28)$$

(The fact that $ab \leq \mu(a)a + \mu^{-1}(b)b$ for all $a, b \geq 0$ follows by separately considering the cases where $\mu(a) \geq b$ and $\mu^{-1}(b) \geq a$.) Choosing $\mu = \phi_2$ in (28) gives

$$\begin{aligned} \Delta_k V_7 &\leq -\frac{\delta}{4(l+1)}k_3(V_1)\Lambda_1(V_1) + \phi_2(k_3(V_1)\Lambda_2(V_1))k_3(V_1)\Lambda_2(V_1) + N_1(x, k)\phi_2(N_1) \\ &\leq -\frac{\delta}{8(l+1)}k_3(V_1)\Lambda_1(V_1) + N_1(x, k)\phi_2(N_1) \quad (\text{by (26)}). \end{aligned} \quad (29)$$

Therefore, since $N_1 \in \mathcal{USB}$ and $\phi_2 \in \mathcal{K}_\infty$, a suitable function $\phi_3 \in \mathcal{K}_\infty$ gives

$$\Delta_k V_7 \leq -\frac{\delta}{8(l+1)}k_3(V_1)\Lambda_1(V_1) + N_1(x, k)\phi_3(V_1). \quad (30)$$

Arguing as in the construction of k_2 above (but with Γ replaced by ϕ_3) provides $k_5 \in \mathcal{K}_\infty$ such that $\Delta_k(k_5 \circ V_1)(x, k) \leq \Delta_k V_1(x, k)\phi_3(V_1(x, k)) \leq -N_1(x, k)\phi_3(V_1(x, k))$. Hence, $V_8 := V_7 + k_5(V_1) \in \mathcal{UPPD}$ satisfies

$$\Delta_k V_8(x, k) \leq -\frac{\delta}{8(l+1)}k_3(V_1(x, k))\Lambda_1(V_1(x, k)) \leq -\alpha_3(|x|), \quad (31)$$

where

$$\alpha_3(s) := \frac{\delta}{8(l+1)} \min\{k_3(u)\Lambda_1(u) : \alpha_1(s) \leq u \leq \alpha_2(s)\}.$$

Since $\alpha_3 \in \mathcal{PD}$, V_8 satisfies the requirements of the theorem. This and Lemma 2 proves the theorem.

5.2 Results for Hybrid Systems

5.2.1 Proof of Theorem 11

For each $k \in \mathbb{Z}_{\geq 0}$, let $V_{\text{cts}}(x, t, k)$ denote the continuous-time strictification of V obtained in [13] for the nonstrictness parameter $q \in \mathcal{P}_{\text{cts}}$. Thus,

$$V_{\text{cts}}(x, t, k) := \left[1 + \frac{1}{\tau} \int_{t-\tau}^t \int_z^t q(\nu) d\nu dz \right] V(x, t, k).$$

The results from [13] show that $\mathcal{D}V_{\text{cts}}(x, t, k) \leq -(\varepsilon/\tau)V_{\text{cts}}(x, t, k)$ for all $x \in C$, $t \geq 0$, and $k \in \mathbb{Z}_{\geq 0}$. We next rewrite the first decay condition in (14) as

$$V(F(x, k), t, k+1) - V(x, t, k) \leq -p(k+1)V(x, t, k), \quad \forall x \in D, t \in [0, \infty), k \in \mathbb{Z}_{\geq 0} \quad (32)$$

where $k \mapsto p(k) := 1 - e^{-r(k)}$ is again of PE type. For each $t \geq 0$, let $V_{\text{dis}}(x, t, k)$ be the strictification

$$V_{\text{dis}}(x, t, k) := \left[1 + \frac{1}{4(l+1)} \sum_{s=k-l}^k \sum_{j=s}^k p(j) \right] V(x, t, k)$$

of V from Theorem 7. The proof of Theorem 7 shows we can take $\kappa(s) \equiv \gamma(s) \equiv s$, and therefore also

$$\Delta_k V_{\text{dis}}(x, t, k) = V_{\text{dis}}(F(x, k), t, k+1) - V_{\text{dis}}(x, t, k) \leq -\frac{\delta}{4(l+1)}V_{\text{dis}}(x, t, k) \quad \forall t \in [0, \infty), k \in \mathbb{Z}_{\geq 0}$$

for all $x \in D$. By enlarging l from the PE assumption as necessary, we can assume $\delta < l$. It follows that the discrete decay condition in (14) holds with V replaced by V_{dis} and with the constant

$$r(k) \equiv \ln \left(\frac{4(l+1)}{4(l+1) - \delta} \right) > 0.$$

Since $\mathcal{D}V \leq 0$ on C and $\Delta_k V \leq 0$ on D , we have $\mathcal{D}V_{\text{dis}} \leq 0$ on C and $\Delta_k V_{\text{cts}} \leq 0$ on D . The uniform boundedness of $S(k)$ and $\int_{t-\tau}^t \int_z^{\bar{z}} q(\nu) d\nu dz$ from Lemma 4 provides constants $r_c, r_d > 0$ such that

$$V_{\text{cts}}(x, t, k) \leq r_c V_{\text{dis}}(x, t, k) \leq r_d V_{\text{cts}}(x, t, k)$$

everywhere. One therefore easily checks that $V^\sharp(x, t, k) := V_{\text{cts}}(x, t, k) + V_{\text{dis}}(x, t, k)$ as given by (15) is an exponential decay Lyapunov function for the hybrid dynamic \mathcal{H} . The periodicity assertion follows as before from Lemma 4, so the result follows from Lemma 6.

5.2.2 Proof of Theorem 14

For each $t \geq 0$ we apply the first part of the proof of Theorem 10 to the functions $(x, k) \mapsto V_1(x, t, k)$ and $(x, k) \mapsto V_2(x, t, k)$ to get V_5 that satisfies

$$\Delta_k V_5(x, t, k) \leq -p(k+1)\Lambda_1(V_1(x, t, k)) + \Lambda_2(V_1(x, t, k))\phi_2(N_1(x, t, k)) \quad \forall x \in D, t \geq 0, k \in \mathbb{Z}_{\geq 0}. \quad (33)$$

This can be done with $\Lambda_1 \in C^1$ and Λ_2 independent of t . For each $k \in \mathbb{Z}_{\geq 0}$, we next apply the continuous time analog of the preceding argument (which is almost exactly the same except with $\Delta_k V_i$ replaced by $\mathcal{D}V_i$ for $i = 1, 2, \dots, 5$, as discussed in the appendix below) to get a continuous version V_5^{cts} of V_5 that satisfies

$$\mathcal{D}V_5^{\text{cts}}(x, t, k) \leq -q(t)\Lambda_1(V_1(x, t, k)) + \Lambda_2(V_1(x, t, k))\phi_2(N_1(x, t, k)) \quad \forall x \in C, t \geq 0, k \in \mathbb{Z}_{\geq 0}. \quad (34)$$

In fact, by enlarging k_2 as necessary (e.g., by enlarging Γ in the discrete version of the proof), we can assume V_5^{cts} and V_5 have the same formula. Applying the strictification method from Theorem 7 to V_5 produces

$$V_6^{\text{dis}}(x, t, k) := V_5(x, t, k) + \frac{1}{4(l+1)}S(k)\Lambda_1(V_5(x, t, k))$$

that satisfies (25) with V_6 replaced by V_6^{dis} and with V_1 and N_1 now also depending on t . Similarly, we apply the continuous time strictification from [13] (as in the proof of Theorem 11) to V_5^{cts} to get

$$V_6^{\text{cts}}(x, t, k) := V_5^{\text{cts}}(x, t, k) + \frac{1}{\tau} \left[\int_{t-\tau}^t \int_z^{\bar{z}} q(\nu) d\nu dz \right] \Lambda_1(V_5^{\text{cts}}(x, t, k))$$

that satisfies $\mathcal{D}V_6^{\text{cts}}(x, t, k) \leq -\Lambda_1(V_1(x, t, k)) + \Lambda_2(V_1(x, t, k))\phi_2(N_1(x, t, k))$ when $x \in C$, possibly by reducing Λ_1 and increasing $\Lambda_2 \in \mathcal{K}_\infty$ without relabeling. Setting $V_6 = V_6^{\text{cts}} + V_6^{\text{dis}}$, and assuming without loss of generality that $1 > \delta/\{4(l+1)\}$ (by enlarging l without relabeling as before), it follows from the fact that V_5^{cts} and V_5 have the same formula that we can enlarge Λ_2 sufficiently so that

$$\Delta_k V_6(x, t, k) \leq -\frac{\delta}{4(l+1)}\Lambda_1(V_1(x, t, k)) + \Lambda_2(V_1(x, t, k))\phi_2(N_1(x, t, k)) \quad \forall x \in D \quad (35)$$

$$\mathcal{D}V_6(x, t, k) \leq -\frac{\delta}{4(l+1)}\Lambda_1(V_1(x, t, k)) + \Lambda_2(V_1(x, t, k))\phi_2(N_1(x, t, k)) \quad \forall x \in C \quad (36)$$

hold for all $t \geq 0$ and $k \in \mathbb{Z}_{\geq 0}$. (This can be seen by bounding $|\Lambda'_1|$ on the relevant intervals and recalling that $F, G \in \mathcal{USB}$ and $V_1 \in \mathcal{UPPD}$. In particular, to get (35), we write $\Delta_k(\Lambda_1 \circ V_5)(x, t, k) = \Lambda'_1(\eta V_5(F(x, k), t, k+1) + (1-\eta)V_5(x, t, k))\Delta_k V_5(x, t, k)$ for $\eta \in [0, 1]$ depending on x, t , and k and use the fact that $\Lambda'_1 \geq 0$ everywhere.) We next follow the remainder of the proof of Theorem 10 applied to V_6 for each $t \geq 0$ to get a function $V_8^{\text{dis}}(x, t, k)$ satisfying the conclusion of the proof when $x \in D$. We also apply the continuous time analog of that part of the proof to V_6 for each k (with $\Delta_k V_i$ replaced by $\mathcal{D}V_i$ for all i as before, similarly to the argument done in the appendix below) to get V_8^{cts} satisfying

$$\mathcal{D}V_8^{\text{cts}}(x, t, k) \leq -\tilde{\alpha}(|x|), \quad \forall x \in C, t \geq 0, k \in \mathbb{Z}_{\geq 0}$$

for a suitable $\tilde{\alpha} \in \mathcal{PD}$. By enlarging $k_4, k_5 \in \mathcal{K}_\infty$ and reducing $k_3 \in \mathcal{PD}$ in the continuous and discrete versions of the proof, we can assume they are the same in both versions, so V_8^{cts} and V_8^{dis} have the same expression. Hence, we can satisfy the requirements of the theorem with their common value. Combined with the result of Lemma 6, this proves the theorem.

6 Examples

One class of systems covered by our discrete time results is as follows. Assume (1) is GAS and that a strict Lyapunov function V for the system is available. This provides $\alpha_1, \alpha_2 \in \mathcal{K}_\infty$ and $\alpha_3 \in \mathcal{PD}$ such that $\Delta_k V(x, k) \leq -\alpha_3(|x|)$ and $\alpha_1(|x|) \leq V(x, k) \leq \alpha_2(|x|)$ everywhere. Assume now that the system is acted on by a PE term $p \in \mathcal{P}_{\text{dis}}$ that freezes the dynamics for certain times. The new system becomes

$$x_{k+1} = [1 - p(k+1)]x_k + p(k+1)F(x_k, k). \quad (37)$$

Thus the new dynamic $F_p(x, k) := [1 - p(k+1)]x + p(k+1)F(x, k)$ fixes the state when $p(k+1) = 0$. By separately considering the cases $p(k+1) = 0$ and $p(k+1) = 1$, one checks that if $p(k) \in \{0, 1\}$ for all k , then $V(F_p(x, k), k+1) - V(x, k) \leq -p(k+1)\alpha_3(|x|) \leq -p(k+1)\Theta(V(x, k))$ everywhere, where $\Theta(s) = \min\{\alpha_3(p) : \alpha_2^{-1}(s) \leq p \leq \alpha_1^{-1}(s)\}$. Since $\Theta \in \mathcal{PD}$, V satisfies the PE decay condition from Theorem 7 for the new dynamic F_p . More generally, assume $p(k) \in [0, 1]$ for all k . Assume also that $V(x, k)$ is a Lyapunov function for (1) that is independent of k and convex in x . Choose $\alpha_3 \in \mathcal{PD}$ such that $V(F(x, k)) - V(x) \leq -\alpha_3(|x|)$ everywhere. Then

$$\begin{aligned} V(F_p(x, k)) - V(x) &\leq [1 - p(k+1)]V(x) + p(k+1)V(F(x, k)) - V(x) \\ &\leq -p(k+1)V(x) + p(k+1)[V(x) - \alpha_3(|x|)] = -p(k+1)\alpha_3(|x|) \end{aligned}$$

everywhere, so F_p again satisfies our PE assumptions.

A general class of *hybrid* systems covered by our strictification results is as follows. Assume the continuous time system (2) admits $q \in \mathcal{P}_{\text{cts}}$, $\gamma \in \mathcal{K}_\infty$, $V \in C^1$, and $\alpha_1, \alpha_2 \in \mathcal{K}_\infty$ satisfying $\mathcal{D}V(x, t) \leq -q(t)\gamma(V(x, t))$ and $\alpha_1(|x|) \leq V(x, t) \leq \alpha_2(|x|)$ for all $x \in \mathbb{R}^n$ and $t \geq 0$ (i.e., (2) admits a nonstrict Lyapunov function in the sense of [13]).¹ Given subsets $C, D \subseteq \mathbb{R}^n$ and $p \in \mathcal{P}_{\text{dis}}$ taking all its values in $\{0, 1\}$, we determine conditions on $F \in \mathcal{USB}$ guaranteeing that we can construct a Lyapunov function for

$$\mathcal{H}_p := \begin{cases} \dot{x} &= G(x, t) \text{ , } & x \in C \\ x_{k+1} &= F_p(x_k, k) \text{ , } & x_k \in D \end{cases} \quad (38)$$

where F_p is as defined above. (The construction we are about to give also works if instead of assuming $p(k) \in \{0, 1\}$ for all $k \in \mathbb{Z}_{\geq 0}$, we assume (i) $x \mapsto V(x, t)$ is convex for each $t \in [0, \infty)$ and (ii) $p(k) \in [0, 1]$ for all $k \in \mathbb{Z}_{\geq 0}$. This situation arises if $\dot{x} = G(x, t) := A(t)x$ is GAS and $A(t)$ is continuous and bounded since then we can take $V(x, t) := x^\top P(t)x$ for a suitable bounded everywhere positive definite matrix $P(t)$ [10, Section 4.6].) To this end, first notice that by reducing $\gamma \in \mathcal{K}_\infty$ as necessary, we can assume $\gamma \in C^1$ and $\gamma(s) \leq \alpha_1(\alpha_2^{-1}(s))/2$ for all $s \geq 0$. Let F satisfy $|F(x, k)| \leq \alpha_2^{-1}(\alpha_1(|x|)/2)$ for all $x \in D$ and $k \in \mathbb{Z}_{\geq 0}$. (This reduces to a linear growth condition when $V(x, t) = x^\top P(t)x$ and P has bounded positive eigenvalues.) By separately considering the cases $p(k+1) = 0$ and $p(k+1) = 1$, it follows that

$$\begin{aligned} V(F_p(x, k), t) - V(x, t) &\leq p(k+1)\alpha_2(|F(x, k)|) - p(k+1)\alpha_1(|x|) \\ &\leq -\frac{1}{2}p(k+1)\alpha_1(|x|) \leq -p(k+1)\gamma(\alpha_2(|x|)) \leq -p(k+1)\gamma(V(x, t)) \end{aligned}$$

for all $x \in D$, $t \geq 0$, and $k \in \mathbb{Z}_{\geq 0}$. A slight variant of the proof of Theorem 11 therefore provides an explicit globally smooth strict Lyapunov function for \mathcal{H}_p having the form

$$V^\sharp(x, t, k) := 2V(x, t) + \frac{1}{\tau} \left[\int_{t-\tau}^t \int_s^t q(r) dr ds \right] \gamma(V(x, t)) + \left[\frac{1}{4(l+1)} \sum_{s=k-l}^k \sum_{j=s}^k p(j) \right] \gamma(V(x, t))$$

for l and τ as in the requirements $p \in \mathcal{P}_{\text{dis}}$ and $q \in \mathcal{P}_{\text{cts}}$ so \mathcal{H}_p is GAS, as claimed.

¹A concrete example where this occurs and where it is easy to find V is where $q \in \mathcal{P}_{\text{cts}}$ (e.g. $q(t) = \sin^2(t)$) and $\dot{x} = h(x, t)$ is GAS (e.g. $\dot{x} = -x$) and we take the dynamic $G(x, t) = q(t)h(x, t)$ and a Lyapunov function $V(x, t)$ for $\dot{x} = h(x, t)$.

7 Conclusions

We provided new methods for constructing closed form strict Lyapunov functions for hybrid systems that admit appropriate nonstrict Lyapunov functions. Our results cover cases where the given nonstrict Lyapunov functions satisfy a decay condition involving persistency of excitation parameters or hybrid versions of the conditions of Matrosov's Theorem. Due to the ubiquity of Lyapunov functions in engineering applications, we expect that our results will be useful in a wide range of settings in which explicit Lyapunov functions are needed such as Lyapunov-based controller design and robustness analysis. We conjecture that our results can be extended to hybrid control systems with outputs. This would extend [25, 26] and the input-to-output stability Lyapunov function constructions from [12] to hybrid systems and also provide more explicit constructions that would be suited for applications.

Appendix

Our proof of Theorem 14 was based on a continuous time version of Theorem 10. We next give a precise statement and proof of this continuous time result, which is applied in the proof of Theorem 14 to $(x, t) \mapsto V_i(x, t, k)$ for each k and $i = 1, 2$. We assume the following version of the Matrosov conditions:

Assumption A.1 *There exist $V_1 : \mathbb{R}^n \times [0, \infty) \rightarrow [0, \infty)$ of class \mathcal{UPPD} and $V_2 : \mathbb{R}^n \times [0, \infty) \rightarrow \mathbb{R}$ of class \mathcal{USB} that are C^1 , $\phi_2 \in \mathcal{K}_\infty$, nonnegative functions $N_1, N_2 \in \mathcal{USB}$, a function $\chi : \mathbb{R}^n \times [0, \infty) \times [0, \infty) \rightarrow \mathbb{R}$, a positive increasing function ϕ_1 , $W \in \mathcal{PD}$, and $q \in \mathcal{P}_{cts}$ such that*

$$\begin{aligned} \mathcal{D}V_1(x, t) &\leq -N_1(x, t), & \mathcal{D}V_2(x, t) &\leq -N_2(x, t) + \chi(x, N_1(x, t), t), \\ |\chi(x, N_1(x, t), t)| &\leq \phi_1(|x|)\phi_2(N_1(x, t)), & \text{and } N_1(x, t) + N_2(x, t) &\geq q(t)W(x) \end{aligned}$$

hold for all $x \in \mathbb{R}^n$ and $t \in [0, \infty)$.

Notice that V_2 can take both positive and negative values. We show:

Theorem A.2 *If (2) satisfies Assumption A.1, then one can construct an explicit strict Lyapunov function for (2). In particular, (2) is GAS.*

To prove this theorem, we indicate the changes needed in the proof of Theorem 10. We define V_3 and λ as in Section 5.1.2 which therefore satisfy

$$\mathcal{D}V_3(x, t) \leq -q(t)W(x) + \phi_1(|x|)\phi_2(N_1(x, t)) \leq -q(t)\lambda(V_1(x, t)) + \phi_1(|x|)\phi_2(N_1(x, t))$$

everywhere. We also define k_1 , Λ_1 , and $V_4 := k_1(V_1)V_3$ as before and as before also determine positive increasing functions Γ and Λ_2 such that

$$\mathcal{D}V_4(x, t) \leq [-\mathcal{D}V_1(x, t)]\Gamma(V_1(x, t)) - q(t)\Lambda_1(V_1(x, t)) + \Lambda_2(V_1(x, t))\phi_2(N_1(x, t)). \quad (\text{A.1})$$

Choosing $k_2 \in \mathcal{K}_\infty$ such that $k_2' \geq \Gamma$ everywhere gives $\mathcal{D}(k_2 \circ V_1) = k_2'(V_1)\mathcal{D}V_1 \leq \Gamma(V_1)\mathcal{D}V_1$, since $\mathcal{D}V_1 \leq 0$ everywhere. Enlarging $k_2 \in \mathcal{K}_\infty$ as necessary, it follows that $V_5 := V_4 + k_2(V_1) \in \mathcal{UPPD}$ satisfies

$$\mathcal{D}V_5(x, t) \leq -q(t)\Lambda_1(V_1(x, t)) + \Lambda_2(V_1(x, t))\phi_2(N_1(x, t)). \quad (\text{A.2})$$

Applying the continuous time strictification method of [13] and enlarging Λ_2 and reducing Λ_1 as necessary without relabeling provides $\gamma \in \mathcal{K}_\infty$ and $\tau > 0$ such that

$$V_6(x, t) := V_5(x, t) + \left[\int_{t-\tau}^t \int_s^t q(r) dr ds \right] \gamma(V_5(x, t)) \quad (\text{A.3})$$

satisfies $\mathcal{D}V_6(x, t) \leq -\Lambda_1(V_1(x, t)) + \Lambda_2(V_1(x, t))\phi_2(N_1(x, t))$. This uses the global boundedness of the double integral in (A.3) from Lemma 4. Arguing as in the proof of Theorem 10 gives $k_3 \in \mathcal{PD} \cap C^1$ such that

$$k_3(r) \leq \phi_2^{-1} \left(\frac{\Lambda_1(r)}{1 + \Lambda_2(r)} \right) \frac{1}{1 + \Lambda_2(r)}, \text{ hence } \phi_2(k_3(V_1)\Lambda_2(V_1)) \Lambda_2(V_1) \leq \Lambda_1(V_1)$$

everywhere. Choose $k_4 \in \mathcal{K}_\infty \cap C^1$ such that $k'_4(s) \geq |k'_3(s)|(\alpha_6 \circ \alpha_1^{-1})(s)$ everywhere, where α_1 is as in the UPPD requirement on V_1 , and $\alpha_6(|x|) \geq V_6(x, t)$ for all $x \in \mathbb{R}^n$ and $t \geq 0$. Then $k'_4(V_1) \geq |k'_3(V_1)|V_6$ everywhere, so $V_7 := k_3(V_1)V_6 + k_4(V_1) \in \mathcal{U}PPD$ everywhere satisfies

$$\begin{aligned} \mathcal{D}V_7 &\leq -k_3(V_1)\Lambda_1(V_1) + k_3(V_1)\Lambda_2(V_1)\phi_2(N_1) - V_6|k'_3(V_1)|\mathcal{D}V_1 + k'_4(V_1)\mathcal{D}V_1 \\ &\leq -k_3(V_1)\Lambda_1(V_1) + k_3(V_1)\Lambda_2(V_1)\phi_2(N_1). \end{aligned}$$

The rest of the argument is similar to the corresponding part of the proof of Theorem 10 with Δ_k replaced by \mathcal{D} and $\frac{\delta}{8(l+1)}$ replaced by 1.

References

- [1] Albertini, F., and E.D. Sontag, “Continuous control-Lyapunov functions for asymptotically controllable time-varying systems,” *International Journal of Control*, **72**(1999), pp. 1630-1641.
- [2] Angeli, D., “Input-to-State Stability of PD-controlled robotic systems,” *Automatica*, **35**(1999), pp. 1285-1290.
- [3] Angeli, D., and E.D. Sontag, “Forward completeness, unboundedness observability, and their Lyapunov characterizations,” *Systems and Control Letters*, **38**(1999), pp. 209-217.
- [4] Angeli, D., E.D. Sontag, and Y. Wang, “A characterization of integral input to state stability,” *IEEE Transactions on Automatic Control*, **45**(2000), pp. 1082-1097.
- [5] Bacciotti, A., and L. Rosier, *Liapunov Functions and Stability in Control Theory*, Springer, London, 2001.
- [6] Cai, C., A. Teel, and R. Goebel, “Converse Lyapunov theorems and robust asymptotic stability for hybrid systems,” in *Proceedings of the 24th American Control Conference (Portland, OR, June 2005)*, pp. 12-17, <http://www.ccec.ece.ucsb.edu/~cai/>.
- [7] Collins, P., “A trajectory-space approach to hybrid systems,” in *Proceedings of the International Symposium on the Mathematical Theory of Networks and Systems (Katholiek Univ. Leuven, Belgium, August 2004)*, Paper #250, <http://homepages.cwi.nl/~collins/>.
- [8] Coron, J.-M., “Global asymptotic stabilization for controllable systems without drift,” *Mathematics of Control, Signals & Systems*, **5**(1992), pp. 295-312.
- [9] Faubourg, L., and J.-B. Pomet, “Control Lyapunov functions for homogeneous “Jurdjevic-Quinn” systems,” *ESAIM: Control, Optimisation and Calculus of Variations*, **5**(2000), pp. 293-311.
- [10] Khalil, H., *Nonlinear Systems, Third Edition*, Prentice Hall, Englewood Cliffs, NJ, 2002.
- [11] Krichman, M., E.D. Sontag, and Y. Wang, “Input-output-to-state stability,” *SIAM Journal on Control and Optimization*, **39**(2001), pp. 1874-1928.
- [12] Malisoff, M., and F. Mazenc, “Further constructions of strict Lyapunov functions for time-varying systems,” in *Proceedings of the American Control Conference (Portland, OR, June 2005)*, pp. 1889-1894.
- [13] Malisoff, M., and F. Mazenc, “Further remarks on strict input-to-state stable Lyapunov functions for time-varying systems,” *Automatica*, **41**(2005), pp. 1973-1978.
- [14] Mancilla-Aguillar, J., R. Garcia, E.D. Sontag, and Y. Wang, “On the representation of switched systems with inputs by perturbed control systems,” *Nonlinear Analysis: Theory, Methods & Applications*, **60**(2005), pp. 1111-1150.
- [15] Mancilla-Aguillar, J., R. Garcia, E.D. Sontag, and Y. Wang, “Uniform stability properties of switched systems with switchings governed by digraphs,” *Nonlinear Analysis: Theory, Methods & Applications*, **63**(2005), pp. 472-490.

- [16] Mazenc, F., “Strict Lyapunov functions for time-varying systems,” *Automatica*, **39**(2003), pp. 349-353.
- [17] Mazenc, F., and M. Malisoff, “Further constructions of control-Lyapunov functions and stabilizing feedbacks for systems satisfying the Jurdjevic-Quinn conditions,” *IEEE Transactions on Automatic Control*, **51**(2006), pp. 360-365.
- [18] Mazenc, F., and D. Nesić, “Lyapunov functions for time varying systems satisfying generalized conditions of Matrosov theorem,” in *Proceedings of the 44th IEEE Conference on Decision & Control (CDC) & European Control Conference ECC 05 (Seville, Spain, December 2005)*, pp. 2432-2437.
- [19] Morin, P., C. Samson, and J.-B. Pomet, “Design of homogeneous time-varying stabilizing control laws for driftless systems via oscillatory approximation of Lie brackets in closed loop,” *SIAM Journal on Control and Optimization*, **38**(1999), pp. 22-49.
- [20] Nesić, D., and A. Loria, “On uniform asymptotic stability of time-varying parameterized discrete-time cascades,” *IEEE Transactions on Automatic Control*, **49**(2004), pp. 875-887.
- [21] Nesić, D., A. Teel, and E.D. Sontag, “Formulas relating \mathcal{KL} stability estimates of discrete-time and sampled-time nonlinear systems,” *Systems and Control Letters*, **38**(1999), pp. 49-60.
- [22] Samson, C., “Velocity and torque feedback control of a nonholonomic cart,” in *Advanced Robot Control (Grenoble, 1990)*, Lecture Notes in Control and Information Sciences Vol. 162, Springer, Berlin, 1991, pp. 125-151.
- [23] Sontag, E.D., “Smooth stabilization implies coprime factorization,” *IEEE Transactions on Automatic Control*, **34**(1989), pp. 435-443.
- [24] Sontag, E.D., “Feedback stabilization of nonlinear systems,” in *Robust Control of Linear Systems and Nonlinear Control* (M.A. Kaashoek, Ed.), Birkhäuser, Basel, 1990, pp. 61-81.
- [25] Sontag, E.D., and Y. Wang, “Notions of input to output stability,” *Systems and Control Letters*, **38**(1999), pp. 235-248.
- [26] Sontag, E.D., and Y. Wang, “Lyapunov characterizations of input to output stability,” *SIAM Journal on Control and Optimization*, **39**(2001), pp. 226-249.
- [27] Van der Schaft, A., and H. Schumacher, *An Introduction to Hybrid Systems*, Lecture Notes in Control and Information Sciences Vol. 251, Springer-Verlag London, Ltd., London, 2000.